



BILATERAL ASYMMETRY IN OTOLITH SIZE OF *Pampus argenteus* (OSTEICHTHYES: STROMATIDAE) FROM IRAQI MARINE WATERS

Audai Qasim¹, Laith A. Jawad^{2*}, Baradi Waryani³

¹ Marine Science Centre, University of Basrah, Basrah, Iraq

² School of Environmental and Animal Sciences, Unitec Institute of Technology, 139 Carrington Road, Mt Albert, Auckland 1025, New Zealand

³ Department of Fresh Water Biology and Fisheries, University of Sindh, Jamshoro, Sindh, Pakistan

*Corresponding Author: laith_jawad@hotmail.com

ARTICLE INFO

Received: 6 February 2022

Accepted: 26 April 2022

Keywords:

Bilateral asymmetry
Otolith
Stromateidae
Pomfret
Basrah

ABSTRACT

Bilateral asymmetry is presumed to reveal the developmental variability of the fish in polluted aquatic environments. In these habitats, high-level asymmetry develops, and these fish expend more energy to balance their growth than fish that are not under an impact. A total of 121 specimens of *Pampus argenteus* were collected from Khor Abdulla located in the northwest part of the Arabian Gulf. The asymmetry of two otolith parameters of the marine fish species *Pampus argenteus*, length and width, was calculated. The results demonstrated that the level of asymmetry was highest for otolith width. The level of asymmetry in both otolith parameters was lowest in fish length ranging between 70-100 mm and the highest in fish ranging between 281-310 mm.

How to Cite

Qasim, A., Jawad, A. L., Waryani, B. (2022): Bilateral asymmetry in otolith size of *Pampus argenteus* (Osteichthyes: Stromatidae) from Iraqi marine waters. Croatian Journal of Fisheries, 80, 103-112. DOI: 10.2478/cjf-2022-0011.

INTRODUCTION

Three pairs of otoliths (asteriscus, lapillus and sagitta) are found inside the inner ear of fish. The sagittae are generally the largest otoliths in most teleosts. Otoliths are composed of calcium carbonate and are involved in hearing and balance functions. Otolith growth is accomplished by the continuous deposition of calcium carbonate layers on an organic matrix. In contrast to bones and scales, otoliths are metabolically passive, and the deposited material remains unchanged and cannot be resorbed (Campana and Neilson, 1985). Otoliths represent important structures that can be used to document life history events in fish (Lecomte-Finiger, 1999).

Otolith shape is genetically determined (L'Abée-Lund, 1988) and is extremely species-specific and therefore useful in studying phylogenetic relationships between species (Lombarte and Leonart, 1993). However, environmental conditions may affect the metabolism of fish, which in turn disturbs somatic growth and accordingly the amount of material deposited in otoliths (Cardinale et al., 2004; Galley et al., 2006; Stransky et al., 2008). Other factors, such as ontogenetic stage as represented by size (Hüssy, 2008), age (Castonguay et al., 1991) and/or sexual maturity status (Mérigot et al., 2007), may also impact the shape of otoliths.

Over the last few decades, the majority of studies exploring the causes of variation in otolith shape have focused on the effect of extrinsic factors, such as environmental conditions, or individual characteristics, such as genotype or state (Mille et al., 2015). However, only a few studies have investigated the possible reasons for intra-individual variation in otolith shape. For example, the difference between otolith shape in the right and left inner ears (referred to as otolith location side). The position of the otolith in the inner ear structure of the fish showed that the two divisions of the vestibular system are exactly the same with three orthogonal semicircular canals that permit perceiving angular accelerations and three otolithic organs devoted to hearing and balance (Panfili et al., 2002). Although there are some interspecific alterations in the size and shape of these features, otoliths are bilaterally symmetrical in roundfishes (Popper and Lu, 2000).

Any changes in the lateralization of the fish body will lead to variation in otolith biomineralization. Accordingly, the deposition of carbonate accretion will be different on both sides of the fish head resulting in asymmetry of otolith mass (Helling et al., 2005), which may induce asymmetry in otolith shape.

Somarakis et al. (1997a) proposed that measurements of fluctuating otolith asymmetry may be a useful indicator of fish condition. Indeed, Clark (1992) revealed that it is a tremendously subtle indicator of stress. The otolith characters examined by Somarakis et al. (1997a) did not show any size or age-related changes. The study of otolith asymmetry is a low-cost, simple measurement

that does not need any special handling or facilities in the field and is not affected by damage during net capture or shrinkage. Fluctuating otolith asymmetry also serves as a potentially useful bioindicator of the health status of different fish populations (Somarakis et al., 1997a, 1997b; Grønckjaer and Sand, 2003), possibly revealing the impact of pollution (Hardersen, 2000), temperature (Lu and Bernatchez, 1999) or parasitism (Sasal and Pampoulie 2000; Reimchen and Nosil, 2001).

In general, the relationship between fish morphology and fluctuating asymmetry has been examined for adult fishes and a number of characters have been surveyed, including the number of gill rakers, pectoral fin rays, fish body proportions, eyespot area, and otolith size and shape (Al-Hassan et al., 1990; Al-Hassan and Hassan, 1994; Somarakis et al., 1997; Jawad, 2001; Øxnevad et al., 2002; Jawad 2003, 2004).

Pampus argenteus is a marine species that prefers to live in the benthopelagic habitats (Riede, 2004) at depths of 5-110 m (Pauly et al., 1996). Individuals of this species are distributed in the Indo-West Pacific region from the Persian Gulf to Indonesia, north of Hokkaido, Japan. They are also reported from the Adriatic (Piper, 2010). This species reaches a maximum total length of 600 mm, with a common length of 300 mm SL (Last, 1997).

Fluctuating asymmetry is usually used to test the health of the environment in which the fish lives (Palmer, 1994). Despite the high commercial value of *P. argenteus*, such a study has never been conducted on this species anywhere within its distributional range to assess the environmental effect on the life of this species.

Assessment of the extent of fluctuating asymmetry has not been performed on the otolith widths or lengths of *Pampus argenteus* examined in the present study. Therefore, the aim of this study is to provide such a study on highly commercially important species like *P. argenteus*.

The aim of the present study was to determine the level of bilateral asymmetry in both the length and width of *Pampus argenteus* otoliths collected from Iraqi marine waters.

MATERIALS AND METHODS

Study area

Khor Abdulla is one of several marine areas of Iraq that includes the estuary of the Shatt AlArab River at the city of Fao, in the Khor alZubair and Um Qasar regions (Fig. 1). The marine biodiversity of Iraq has gone through several changes related to its geological history, its temporal and geographical positions at the top of the Arabian Gulf, and its physiographical complexity (Jawad, 2016). The climate and hydrology of the lower Mesopotamian plain including its coastal area were fully described by Purser et al. (1982) and AlAzzawi (1986).




Fig 1. Map showing the sampling area of *Pampus argenteus*

Fish specimens

Specimens of *P. argenteus* (121) (Fig. 2) were collected from small commercial trawlers (21 m length x 3.5 m width), fishing with nets of mesh size 2.5 cm off Khor Abdullah at the southern extent of Iraqi marine waters. The sampling area was chosen as it represents one of the main fishing grounds for the species in Iraq. The specimens were caught in the period February-September 2019 at depths of 10-25 m. Sagittae from both sides of the fish head were extracted from the sacculus part of the fish inner ear. The standard length (SL) of the fish samples ranged from 78 to 293 mm. Otolith length and width were measured to the nearest millimetre under a dissecting microscope (Fig. 3). The features chosen for bilateral asymmetry analysis were previously used in fish studies (Al-Rasady et al., 2010; El-Regal et al., 2016).

Data analysis

Statistical analysis was based on the squared coefficient of asymmetry variation (CV_a^2) for the two otolith dimensions according to Valentine et al. (1973):

$$CV_a^2 = (S_{r-1} \times 100 / X_{r+1})^2$$

where S_{r-1} is the standard deviation of signed differences and X_{r+1} is the mean of the character, which was calculated by adding the absolute scores for both sides and dividing by the sample size. Bilateral asymmetry values and measurement errors were small and normally distributed around a mean of zero (Merilä and Bjöklund,

1995). Individual inaccuracies in taking measurements can disturb the results of bilateral asymmetry analysis, rendering it vague (Palmer, 1994). Therefore, in the present study all measurements were carried out in duplicate by only one person in order to reduce any unsolicited errors (Lee, 1990). Coefficients of asymmetry were compared between the different SL classes using ANOVA.

RESULTS

The results of the asymmetry analysis of the data of otolith length and width are summarized in Table 1. The results indicated that the level of asymmetry was highest for otolith width.

Table 1. Squared coefficient of asymmetry (CV_a^2) value and character means (X_{r+1}) of *P. argenteus* collected from the marine waters of Iraq

Character	CV_a^2	N	Character mean \pm SD	% of individuals with asymmetry
Otolith length	41.55	121	7.93 \pm 3.1	80
Otolith width	89.56	121	3.86 \pm 2.3	99

Furthermore, the lowest and highest levels of asymmetry were recorded in fish ranging in length between 70-100 mm and 281-310 mm, respectively (Table 2).




Fig2. Specimen of *Pampus argenteus*, 260 mm SL, collected from the marine waters of Iraq




Fig 3. Otolith of *Pampus argenteus*, 326 mm TL, showing the length and width of the otolith. Scale bar = 10 mm.

Table 2. Squared coefficient of asymmetry and character means by size class of *P. argenteus* collected from the marine waters of Iraq

Character	CV _a ²	N	Character mean ± SD	% of individuals with asymmetry
Otolith length				
Fish standard length mm				
70-100	24.6	4	3.34 ± 1.2	100
101-130	29.9	10	5.23 ± 2.2	100
131-160	30.3	9	6.22 ± 2.4	79
161-190	35.7	20	10.93 ± 1.3	98
191-220	39.9	10	11.98 ± 1.6	89
221-250	42.6	49	11.41 ± 1.9	99
251-280	48.7	10	12.74 ± 1.3	100
281-310	50.9	8	12.56 ± 2.5	94
Otolith width				
Fish standard length mm				
70-100	18.1	4	2.1	99
101-130	19.9	10	2.5	100
131-160	19.9	9	3.1	79
161-190	21.9	20	3.7	99
191-220	22.3	10	4.1	100
221-250	22.9	49	4.8	99
251-280	23.1	10	5.2	97
281-310	23.5	8	5.3	98

DISCUSSION

Due to the deficiency of data regarding natural otolith asymmetry in this part of the world, it is difficult to evaluate whether or not they are typical for *P. argenteus*. Although it is possible that the relatively high level of asymmetry in otolith width may be related to environmental factors, it is not possible to support this hypothesis at this stage. The relatively small level of asymmetry observed in otolith length suggests that this parameter may be less vulnerable to environmental impact. In a previous study, Jawad (2003) suggested that the lower level of bilateral asymmetry observed in otolith length of sparid fishes off Libya may be due to the fact that the developmental period of these features may not agree with the existence of opposing environmental events. Palmer and Strobeck (1992) noted that slight changes throughout the growth of the fish can result in a move from normal developmental trails. These irregularities may be due to the state and quantity of

food, excessive temperatures, parasites, disease and/or behavioral burdens imposed by interactions with related species living in the same environment (Markov, 1995). The drawback of measurement error in fluctuating asymmetry (FA) studies is not immaterial but has been given increasing consideration over the last few decades (Palmer and Strobeck, 1986). Since variations in values of bilateral characters are frequently small [$<1-50/0$ of the total variation in a given trait (Palmer, 1996)] and are by explanation random, FA indices are likely to be especially disposed to measurement error. It has been observed that measurement error in fluctuating asymmetry approximations can be high even when the measurement error of the original traits is very low. For example, Merilä and Bjöklund (1995) reported that measuring error in skeletal characters of greenfinches *Curduelis* sp. was 2.5 and 4.5% for the left and the right sides, respectively, while measuring error in fluctuating asymmetry was 98%.

The reason for this is that a large amount of the variation due to measuring error can occur as a result of either high levels of absolute measuring error or low levels of absolute asymmetry (Somarakis et al., 1997a).

Although there is no reference range for bilateral asymmetry levels in otolith dimensions to check the acuteness level of asymmetry, it is possible to assess the observed levels of otolith asymmetry dimensions of *P. argenteus* in the present study by comparing them with available published data on the same species studied in other countries and other fish species. The results obtained in the present study were compared with the values of fluctuating otolith asymmetry sizes in some fish species collected from the same area (Arabian Gulf), from neighbouring areas (Sea of Oman, Red Sea) and other areas (Black Sea) (Table 3).

Fluctuating asymmetry values in otolith length in the equated species ranged between 2.1 in *Sardinella sindensis* collected from the Arabian Gulf (Jawad et al., 2012d) and 88.71 in *Rastrelliger kanagurta* collected off the Omani coast of the Sea of Oman (Al-Mamry et al., 2011a). Fluctuating asymmetry in otolith width ranged

between 4.59 in *Sargocentron spiniferum* off the Egyptian coast of the Red Sea (El-Mahdy et al., 2019) and 117.60 in *Rhynchorhamphus georgi* collected off the Omani coast of the Sea of Oman (Al-Rasady et al., 2010). The fluctuating asymmetry in otolith length in *P. argenteus* was 31.55, close to the mid-point of the maximum value of 88.71 obtained for *Rastrelliger kanagurta* collected off the Omani coast of the Sea of Oman (Al-Mamry et al., 2011a). For otolith width, the level of the fluctuating asymmetry in *P. argenteus* was 89.56, close to the highest value of fluctuating asymmetry (117.60) obtained for *Rhynchorhamphus georgi* collected off the Omani coast of the Sea of Oman (Al-Rasady et al., 2010). The comparative assessment indicates that the degree of asymmetry in both otolith length and width in *P. argenteus* is high and that otolith width is appreciably higher. While the species of fish included in the current comparison are not the same and are from diverse localities, they indicate some degree of environmental impact on otolith features, i.e. characters of the otolith that are either susceptible to or endure unsuitable environmental settings (Fey and Hare, 2008).

Table 3. Comparison of the Coefficient of asymmetry (CV_a^2) of otolith sizes of *P. argenteus* examined in the present study with those of other fish species collected from neighbouring localities

Species	Coefficient of asymmetry (CV_a^2)		Reference
	OL	OW	
<i>Pampus argenteus</i>	41.55	89.56	Present study
<i>Acanthopagrus arabicus</i>	47.74	88.65	Abdulsamad et al. (2020)
<i>Acanthopagrus latus</i>	44.98	88.65	Abdulsamad et al. (2020)
<i>Beryx splendens</i>	41.87	87.30	Al-Busaidi et al. (2010)
<i>Carangoides caeruleopinnatus</i>	28.43	54.05	Jawad et al. (2012a)
<i>Chlorurus sordidus</i>	14.05	10.44	El-Regal et al. (2016)
<i>Hipposcarus harid</i>	15.19	11.90	El-Regal et al. (2016)
<i>Liza Kluzingeri</i>	4.23	14.06	Sadighzadeh et al. (2011)
<i>Lutjanus bengalensis</i>	5.06	10.29	Jawad (2012b)
<i>Merlangius merlangus</i>	4.710	4.772	Kontaş et al. (2018)
<i>Rastrelliger kanagurta</i>	88.71	41.75	Al-Mamry et al. (2011a)
<i>Rhynchorhamphus georgi</i>	66.70	117.60	Al-Rasady et al. (2010)
<i>Sargocentron spiniferum</i>	2.34	4.59	El-Mahdy et al. (2019)
<i>Sardinella sindensis</i>	2.10	9.00	Jawad et al. (2012d)
<i>Sillago sihama</i>	2.9	21.6	Jawad et al. (2012d)
<i>Sparidentex hasta</i>	41.87	87.30	Abdulsamad et al. (2020)
<i>Engraulis australis</i>	43.55	86.56	Jawad and Adams (2021)
<i>Otolithes ruber</i>	39.65	87.53	Jawad et al. (2021)

The sources and reasons for otolith asymmetry in fishes may be due to numerous factors. Although genetic factors might be responsible, they cannot be considered at this stage due to the lack of genetic data on the marine ichthyofauna of Iraq. Environmental factors might also lead to increased levels of asymmetry but may occur at low levels before producing widespread death (Bengtsson and Hindberg, 1985).

Pollution of seawater and marine sediments by hydrocarbons, heavy metals, pesticides and organic matter is considered the chief source of environmental impacts, particularly in the marine waters of Iraq where various pollutants have been noted over the last twenty years (Al-Imarah et al., 2007; Zuhkair et al., 2007; Al-Jaberi and Al-Dabbas, 2014). Chemical and organic pollution can lead to morphological deformities in fish (Elie and Girard, 2014). Indeed, fish deformities in Iraqi marine waters have been associated with heavy metal and organic pollution (Jawad and Bannai, 2014; Jawad et al. 2014, 2017). Other aquatic organisms within Iraqi waters have also been critically distressed due to environmental pollutants (Saeed et al., 1999; Zuhkair et al., 2008). Environmental impacts may also be caused by natural events (Bengtsson and Hindberg, 1985), and it is feasible that in some cases they may also be impacting Iraqi marine waters.

Some authors have demonstrated a connection between the coefficient of asymmetry and fish length (Al-Hassan et al., 1990; Al-Hassan and Hassan, 1994; Al-Hassan and Shwafi, 1997; Jawad, 2001), characterized by increasing levels of asymmetry with fish length (Al-Mamry et al., 2011a, b, c; Jawad et al., 2012a, 2012b, 2012c; Mabrouk et al., 2014). In the current study, it was noted that the larger specimens of *P. argenteus* had higher levels of bilateral asymmetry than smaller specimens ($P < 0.001$). It is possible that the differences may be related to fish growth. Valentine et al. (1973) noted that trait means were always lower in smaller size classes, which may be related to ontogenetic variation and a subsequent increase in bilateral asymmetry with size (age), and/or possible historical events which result in a secular increase in bilateral asymmetry. Thiam (2004) suggested that increasing levels of bilateral asymmetry with fish size could be due to the fact that the larger individuals had longer periods of contact with unfavourable environmental scenarios.

A management policy is urgently required in order to reinstate a healthy environment in the marine waters of Iraq.

ACKNOWLEDGEMENTS

We would like to thank Declan Quigley, Sea-Fisheries Protection Authority, Dublin, Ireland for reading the manuscript. The study was completed without external funding.

BILATERALNA ASIMETRIJA VELIČINE OTOLITA KOD VRSTE *Pampus argenteus* (OSTEICHTHYES: STROMATIDAE) IZ IRAČKIH MORSKIH VODA

SAŽETAK

Pretpostavlja se da bilateralna asimetrija otkriva varijabilnost u razvoju riba u zagađenom vodenom okruženju. U tim se staništima razvija asimetrija visoke razine i te ribe troše više energije da uravnoteže svoj rast od riba koje nisu pod takvim utjecajima. Ukupno 121 primjerak vrste *Pampus argenteus* prikupljen je iz Khor Abdulle koji se nalazi u sjeverozapadnom dijelu Arapskog zaljeva. Izračunata je asimetrija dvaju parametara otolita morske vrste *Pampus argenteus*, duljine i širine. Rezultati su pokazali da je razina asimetrije najveća za širinu otolita. Razina asimetrije u oba parametra otolita najniža je pri duljini riba u rasponu od 70-100 mm, a najviša kod riba u rasponu između 281-310 mm.

Ključne riječi: Bilateralna asimetrija, otolit, Stromateidae, pomfret, Basrah

REFERENCES

- Abdulsamad, S.M., Jawad, L. A., Al-Nusear, A.N., Waryani, B., Rutkayová, J. (2020): Asymmetry in the otolith length and width of three sparid fish species collected from Iraqi waters. *Marine Pollution Bulletin*, 156, p. 111177.
- AlAzzawi, M. (1986): La sédimentation actuelle sur la plaine de la Basse Mésopotamie (Irak). Unpublished thesis, ParisSud Univ. (Orsay), 832: 172 fig., 68 pl.
- Albusaidi, H. K., Jawad, L. A., Al-Mamry, J.M., Al-Marzouqi, M.S., 2010. Fluctuating asymmetry in the otolith width and length of adult teleost (*Beryx splendens* Lowe, 1834) (family: Bercidae) collected from the Arabian Sea coasts of Sultanate of Oman. *Croatian Journal of Fisheries: Ribarstvo*, 68, 159–166.
- Al-Hassan, L. A. J., Hassan, S. S. (1994): Asymmetry study in *Mystus pelusius* collected from Shatt al-Arab River, Basrah, Iraq. *Pakistan Journal of Zoology*, 26, 276-278.
- Al-Hassan, L. A. J., Shwafi, N. A. A. (1997): Asymmetry analysis in two marine teleost fishes collected from the Red Sea coast of Yemen. *Oceanographic Literature Review*, 10, 44, 1205.
- Al-Hassan, L. A. J., Al-Dubaikel, A.Y., Wahab, N. K., Al-Daham, N. K. (1990): Asymmetry analysis in the catfish, *Heteropneustes fossilis* collected from Shatt Al-Arab River, Basrah, Iraq. *Rivista di idrobiologia*, 29, 3, 775-780.
- Al-Imarah, F. J., Hantoosh, A. A., Nasir, A. M. (2007): Petroleum hydrocarbons in water and sediments of northwest Arabian Gulf 1980–2005. *Aquatic Ecosystem Health & Management*, 10, 3, 335-340.

- Al-Jaberi, M. H., Al-Dabbas, M. A. (2014): Assessment of heavy metals pollution in the sediments of Iraqi coastlines. *Science*, 3, 9.
- Al-Saad, H. T., Abd, I. A., Al-Hello, M.A., Zukhair, M. K. (2006): Environmental Assessment of trace metals pollution in sediment of Khor al-Zubair, Iraq. *Marina Mesopotamica*, 21, 2, 23-33.
- Al-Mamry, J.M., Jawad, L., Ambuali, A. (2011a): Fluctuating asymmetry in the otolith length and width of adult Indian mackerel *Rastrelliger kanagurta* (Cuvier, 1817) collected from Muscat waters at the sea of Oman. *Journal of Black Sea/Mediterranean Environment*, 17, 254–259.
- Al-Mamry, J.M., Jawad, L. A., Al-Bimani, S. M. H., Al-Busaidi, H. K., Al-Marzouqi, M. S., Al-Habsi, S. H. (2011b): Asymmetry analysis study on *Callionymus margaretae* Regan, 1906 collected from the Arabian Sea coasts of Oman. *Croatian Journal of Fisheries: Ribarstvo*, 69, 1, 9-9.
- Al-Mamry, J., Jawad, L., Al-Bimani, S., Al-Ghafari, F., Al-Mamry, D., Al-Marzouqi, M. (2011c): Asymmetry in some morphological characters of *Leiognathus equulus* (Forsskål) (Leiognathidae) collected from the Sea of Oman. *Fisheries & Aquatic Life*, 19, 1, 51-55.
- Al-Rasady, I. H., Jawad, L.A., Al-Mamry, J.M., Al-Mamari, H.M., Al-Yarubi, M.M., Al-Mamary, D.S. (2010): Fluctuating asymmetry in the otolith length and width of *Rhynchorhamphus georgi* (Valenciennes, 1846) (Family: Hemiramphidae) collected from the Sea of Oman. *Annals de Museo civico de St. naturale de Ferrara*, 13, 85-89.
- Bengtsson, B. E., Hindberg, M. (1985): Fish deformities and pollution in some Swedish waters. *Ambio*, 14, 32–35.
- Campana, S. E., Neilson, J. D. (1985): Microstructure of fish otoliths. *Canadian Journal of Fisheries and Aquatic Sciences*, 42, 1014–1032.
- Cardinale, M., Doering-Arjes, P., Kastowsky, M., Mosegaard, H. (2004): Effects of sex, stock, and environment on the shape of known-age Atlantic cod (*Gadus morhua*) otoliths. *Canadian Journal of Fisheries and Aquatic Sciences*, 61, 2, 158-167.
- Castonguay, M., Simard, P., Gagnon, P. (1991): Usefulness of Fourier analysis of otolith shape for Atlantic mackerel (*Scomber scombrus*) stock discrimination. *Canadian Journal of Fisheries and Aquatic Science*, 48, 296–302.
- Clark, G. M. (1992): Fluctuating asymmetry: a technique for measuring developmental stress of genetic and environmental origin. *Acta Zoologica Fennica*, 191, 31-36.
- Elie, P., Girard, P. (2014): La santé des poissons sauvages: les codes pathologie, un outil d'évaluation. Association Santé Poissons Sauvages. Achevé d'imprimer sur les Presses d'AVL diffusion, Montpellier (France) (Pages: 286 pp.).
- El-Mahdy, S.M., Osman, Y. A., Mohammad, A.S., Mehanna, S.F. (2019): Fluctuating asymmetry in otolith morphology of sabre squirrelfish, *Sargocentron spiniferum* (Forsskål, 1775) from the red sea. *Egypt. International Journal of Fisheries and Aquatic Studies*, 7, 188–191.
- El-Regal, M.A., Jawad, L.A., Mehanna, S., Ahmad, Y. (2016): Fluctuating Asymmetry in the Otolith of two Parrotfish Species, *Chlorurus sordidus* (Forsskal, 1775) and *Hipposcarus harid* (Forsskal, 1775) from Hurghada, Red Sea coast of Egypt. *International Journal of Marine Science*, 6, 1-5.
- Escos, J., Alados, C.L., Emlen, J.M., Alderstein, S., 1995. Developmental instability in the Pacific Hake parasitized by myxosporeans *Kudoa* spp. *Transactions of the American Fisheries Society*, 124, 6, 943-945.
- Fey, D.P., Hare, J.A. (2008): Fluctuating asymmetry in the otoliths of larval Atlantic menhaden *Brevoortia tyrannus* (Latrobe) — a condition indicator? *Journal of Fish Biology*, 72, 121–130.
- Galley, E. A., Wright, P. J., Gibb, F. M. (2006): Combined methods of otolith shape analysis improve identification of spawning areas of Atlantic cod. *ICES Journal of Marine Science*, 63, 1710–1717.
- Gonçalves, D. M., Simões, P. C., Chumbinho, A. C., Correia, M. J., Fagundes Oliveira, R. F. (2002): Fluctuating asymmetry and reproduction success in the peacock blenny. *Journal of Fish Biology*, 60, 810-820.
- Grønkrjaer, P., Sand, M. K. (2003): Fluctuating asymmetry and nutritional condition of Baltic cod (*Gadus morhua*) larvae. *Marine Biology*, 143, 191–197.
- Hardersen, S. (2000): The role of behavioural ecology of damselflies in the use of fluctuating asymmetry as a bioindicator of water pollution. *Ecological Entomology*, 25, 45–53.
- Helling, K., Scherer, H., Hausmann, S., Clarke, A. H. (2005): Otolith mass asymmetries in the utricle and saccule of flatfish. *Journal of Vestibular Research*, 15, 59–64.
- Hüssy, K. (2008): Otolith shape in juvenile cod (*Gadus morhua*): ontogenetic and environmental effects. *Journal of Experimental Marine Biology and Ecology*, 364, 35–41.
- Jawad, L. A. (2001): Preliminary asymmetry analysis of some morphological characters of *Tilapia zilli* (Pisces: Cichlidae) collected from three localities in Libya. *Bollettino-Museo Regionale Di Scienze Naturali*, 18, 1, 251-257.
- Jawad, L. A. (2003): Asymmetry in some morphological characters of four sparid fishes from Benghazi, Libya. *Oceanological and Hydrobiological Studies*, 32, 83–88.
- Jawad, L. A. (2004): Asymmetry analysis in the mullet, *Liza abu* collected from Shatt al-Arab River, Basrah, Iraq. *Bollettino del Museo Regionale di Scienze Naturali di Torino*, 21, 1, 145-150.
- Jawad, L. A. (2016): Coastal Area of Iraq: An Appraisal of the Fish Biodiversity, Functioning, Threats and Conservation Management. In: Coastal fishes, habitat, behavior and conservation. Martinez, E. K. (editor). Nova Science Publisher, New York, USA, 385pp.

- Jawad, L. A., Adams, N.J. (2021): Fluctuating asymmetry in the size of the otolith of *Engraulis australis* (Shaw, 1790) recovered from the food of the Australasian gannet, *Morus serrator*, Hauraki Gulf, New Zealand. *Marine Pollution Bulletin*, 168, 112391.
- Jawad, L. A., Bannai, M. (2014): Characterization of hyperostosis in *Platax teira* (Forsskål, 1775) collected from marine water of Iraq, North West Arabian Gulf. *Sky Journal of Agriculture Research*, 3: 109-111.
- Jawad, L. A., Al-Faisal, A. J., Al-Mutlak, F. M. (2014): Incidence of Lordosis in the Cyprinid Fish, *Carasobarbus luteus* and the Shad, *Tenuulosa ilisha* Collected from Basrah Waters, Iraq. *International Journal of Marine Science*, 45, 1-5.
- Jawad, L. A., Al-Mamry, J., Al-Mamari, D. (2012a): Fluctuating asymmetry in the otolith width of *Carangoides caeruleopinnatus* (Rüppell 1830) (Carangidae) Collected from Muscat City coast on the sea of Oman. *Croatian Journal of Fisheries*, 70, 125-133.
- Jawad, L. A., Al-Mamry, J. M., Al-Mamari, D. (2012b): Fluctuating asymmetry in the otolith dimensions of *Lutjanus bengalensis* (Lutjanidae) collected from Muscat coast on the sea of Oman. *Biological Journal of Armenia* 64, 117-121.
- Jawad, L. A., Al-Mamry, J. M., Al-Shuaily, S. (2012c): Bilateral asymmetry in some morphological characters of *Parapercis alboguttata* (Günther, 1872) (Family: Pigiipedidae) collected from the Arabian Sea coasts of Oman. *Romanian Journal of Biology and Zoology*, 57, 51-62.
- Jawad, L. A., Qasim, A.M., Al-Faiz, N. A. (2021): Bilateral asymmetry in size of otolith of *Otolithes ruber* (Bloch & Schneider, 1801) collected from the marine waters of Iraq. *Marine Pollution Bulletin*, 165, 112110.
- Jawad, L., Sadighzadeh, Z., Al-Mamary, D. (2012d): Fluctuating asymmetry in the otolith length, width and thickness in two pelagic fish species collected from the Persian Gulf near Bandar Abbas. *Annales: Series Historia Naturalis*, 22, 1, 83.
- Jawad, L., Al-Mamry, J., Al-Bimani, S., Al-Ghafri, F., Al-Marzouqi, M. (2012e): Asymmetry of some morphological characters of *Upeneus doriae* (Osteichthyes: Mullidae) collected from the sea of Oman. *Thalassia Salentina*, 34, 3-10.
- Jawad, L., Al-Mamry, J., Al-Busaidi, J., Al-Mamari, A., Al-Mamry, S., Al-Owisi, K., Al-Rubiey, M., 2012f. Asymmetry in some morphological characters of Indian oil sardine, *Sardinella Longiceps* Valenciennes, 1847 collected from Muscat waters on the Sea of Oman. *Water Research Management*, 2, 61-64.
- Jawad, L. A., Jayabalan, N., Jawad, H. L. (2017): Unilateral microphthalmia and slow growth rate in a natural population of the marine teleost *Pampus argenteus* (Actinopterygii: Perciformes: Stromateidae). *Bulletin of Fish Biology*, 17, 39-44.
- Kontas, S., Bostanci D., Yedier S., Kurucu, G., Polat, N. (2018): Investigation of fluctuating asymmetry in the four otolith characters of *Merlangius merlangus* (Linnaeus, 1758). Collected from Middle Black Sea. *Türk Denizcilik ve Deniz Bilimleri Dergisi*, 4, 128-138.
- Last, P.R. (1997): Stromateidae. Butterfishes, silver pomfrets. In K.E. Carpenter and V. Niem (eds.) *FAO Identification Guide for Fishery Purposes. The Western Central Pacific. The living marine resources of the Western Central Pacific. Volume 4. Bony fishes part 2 (Mugilidae to Carangidae)*. Rome, FAO. 1999. pp. 2069-2790.
- L'Abée-Lund, J. H. (1988): Otolith shape discriminates between juvenile Atlantic salmon, *Salmo salar* L., and brown trout, *Salmo trutta* L. *Journal of Fish Biology*, 33, 899-903
- Lecomte-Finiger, R. (1999): L'otolithe: la «boîte noire» des Téléostéens. *L'Année Biologique*, 38, 107-122.
- Lee, D. H., Lysak, R. L. (1990): Effects of azimuthal asymmetry on ULF waves in the dipole magnetosphere. *Geoph. Research letters*, 17, 53-56.
- Lombarte, A., Leonart, J. (1993): Otolith size changes related with body growth, habitat depth and temperature. *Environmental Biology of Fishes*, 37, 297-306.
- Lu, G. Q., Bernatchez, L. (1999): A study of fluctuating asymmetry in hybrids of dwarf and normal lake whitefish ecotypes (*Coregonus clupeaformis*) from different glacial races. *Heredity*, 83, 742-747.
- Mabrouk, L., Guarred, T., Hamza, A., Hellal, A.N. (2014): Fluctuating asymmetry in grass goby, *Zosterisessor ophiocephalus* Pallas, 1811 inhabiting polluted and unpolluted area in Tunisia. *Marine Pollution Bulletin*, 85, 248-251.
- Markov, T. A. (1995): Evolutionary ecology and developmental instability. *Annual Review of Entomology*, 40, 105-120.
- Mérigot, B., Letourneur, Y., Lecomte-Finiger, R. (2007): Characterization of local populations of the common sole *Solea solea* (Pisces, Soleidae) in the NW Mediterranean through otolith morphometrics and shape analysis. *Marine Biology*, 151, 997-1008.
- Merilä, J., Bjöklund, M. (1995): Fluctuating asymmetry and measurement error. *Systematic Biology*, 44, 97-101.
- Mille, T., Mahe, K., Villanueva, M.C., De Pontual, H., Ernande, B. (2015): Sagittal otolith morphogenesis asymmetry in marine fishes. *Journal of Fish Biology*, 87, 3, 646-663.
- Øxnevad, S. A., Heibo, E., Vollestad, L. A. (2002): Is there a relationship between fluctuating asymmetry and reproductive investment in perch (*Perca fluviatilis*)? *Canadian Journal of Zoology*, 80, 120-125.
- Palmer, A. R. (1994): Fluctuating asymmetry analyses: a premier. In: Markov, T. A. (Ed.), *Developmental Instability: Its Origin and Evolutionary Implications*. Kluwer, Dordrecht, Netherlands, 335-364.

- Palmer, A. R. (1996): Waltzing with asymmetry. *Bioscience*, 46, 518-532.
- Palmer, A. R., Strobeck, C. (1986): Fluctuating asymmetry: measurement, analysis, patterns- *Annual Review of Ecology and Systematics*, 17, 391-421.
- Palmer, A. R., Strobeck, C. (1992): Fluctuating asymmetry as a measure of developmental stability: implications of non-normal distribution and power of stability tests. *Acta Zoologica Fennica*, 191, 57-72.
- Panfili, J., de Pontual, H., Troadec, H., Meunier, F. J. (2002): *Manuel de sclérochronologie des poissons*. Montpellier: Coédition Ifremer-IRD.
- Popper, A. N., Lu, Z. (2000): Structure-function relationships in fish otolith organs. *Fisheries Research*, 46, 15-25.
- Pauly, D., A. Cabanban and F.S.B. Torres Jr. (1996): Fishery biology of 40 trawl-caught teleosts of western Indonesia. p. 135-216. In D. Pauly, P. Martosubroto (eds.) *Baseline studies of biodiversity: the fish resource of western Indonesia*. ICLARM Studies and Reviews 23.
- Piper, R. (2010): Re-occurrence of silver pomfret *Pampus argenteus* in the North Sea. *Marine Biodiversity Record*, 3, e102.
- Purser, B. H., AlAzzawi, M., AlHassani, N. H. et al. 1982. Caractères et évolution du complexe deltaïque Tigre-Euphrate. *Mémoire Society Geology of France*, 144, 207216.
- Riede, K. (2004): Global register of migratory species - from global to regional scales. Final Report of the R&D-Projekt 808 05 081. Federal Agency for Nature Conservation, Bonn, Germany. 329 p.
- Reimchen, T. E., Nosil, P. (2001): Lateral plate asymmetry, diet and parasitism in threespine stickleback. *Journal of Evolutionary Biology*, 14, 632-645.
- Sadighzadeh, Z., Jawad, L., Al-Marzouqi, M. (2011): Fluctuating asymmetry in the otolith of the mugilid fish *Liza kluzingeri* (Day, 1888) from Persian Gulf near Bandar Abbas. *Thalassia Salentina*, 33, 95-102.
- Saeed, T., Al-Ghadban, A.N., Al-Shemmari, H., Al-Mutairi, M., Al-Hashash, H. (1999): Preliminary assessment of the impact of draining of Iraqi marshes on Kuwait's northern marine environment. Part II. Sediment associated pollutants. *Water Science and Technology*, 40, 7, 89-98.
- Sasal, P., Pampoulie, C. (2000): Asymmetry, reproductive success and parasitism of *Pomatoschistus microps* in a French lagoon. *Journal of Fish Biology*, 57, 382-390.
- Somarakis, S., Kostikas, I., Tsimenides, N. (1997a): Fluctuating asymmetry in the otoliths of larval fish as an indicator of condition: conceptual and methodological aspects. *Journal of Fish Biology, Supplement A* 51, 30-38.
- Somarakis, S., Kostikas, I., Peristeraki, N., Tsimenides, N. (1997b): Fluctuating asymmetry in the otoliths of larval anchovy *Engraulis encrasicolus* and the use of developmental instability as an indicator of condition in larval fish. *Marine Ecology Progress Series*, 151, 191-203.
- Stransky, C., Baumann, H., Fevolden, S.E., Harbitz, A., Høie, H., Nedreaas, K. H., Salberg, A.B., Skarstein, T. H. (2008): Separation of Norwegian coastal cod and Northeast Arctic cod by outer otolith shape analysis. *Fisheries Research*, 90, 1-3, 26-35.
- Thiam, N. (2004): Ecomorphologie de *Trisopterus luscus* (Linnaeus, 1758) tacaud, adaptation a la température at l'asymétric fluctuante (Ph.D. Thesis) Faculté des Sciences, Université de Vigo (190 pp.).
- Valentine, D. W., Soule, M. E., Samollow, P. (1973): Asymmetry in fishes: a possible statistical indicator of environmental stress. *Fishery Bulletin*, 71, 357-370.
- Zuhkair, M. K., Abd, I. M., DouAbul, A. A. (2008): Uptake-release of pollutant by hilsa shad *Tenualosa ilisha* (Hamilton-Buchanan) fish collected from southern Iraq. *Mesopotamian Journal of Marine Science*, 23, 237-255.